



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

but we may confidently expect the beginnings of such a genetic psychology in the future. At any rate, in this field, as in most other fields, progress and profit are increased by greater exactness and care, by more accurate and convenient apparatus and by shorter and more definite methods. These elements are the ones which experimental psychology is trying to introduce into the exploration of mental life. The fact that these methods are somewhat new in psychological work gives us the right to call a system of them a 'new psychology.'

Professor Stanley's claim that biology is the main standpoint of psychology is quite justified—if 'psychology' means the science of mental development. It must be remembered, however, that there is a fundamental difference in aim and method which marks off experimental psychology from the other mental sciences. Its object is to determine the fundamental laws of mental activity in the adult human being under ordinary circumstances. The change of the problem to child-study, to the development of the individual or of the race, or to abnormal circumstances, produces closely related sciences. All these sciences are inter-dependent. In fact, all these sciences—as Professor Stanley implies—are needed for a concrete, practical understanding of mental life; nevertheless convenience and clearness sometimes require that attention should be concentrated on one of them at a time.

E. W. SCRIPTURE.

NEW HAVEN, CONN., May 20, 1898.

#### FOSSIL FULGUR PERVERSUM AT AVALON, N. J.

ON page 682 of SCIENCE the quotation from Captain Swain, of the Avalon Life Saving Station, N. J., with reference to the casting ashore of *Fulgur perversum* is slightly inaccurate. I now quote from his letter the passage I read at the Academy that "the conchs in question come ashore only during a strong northwest (not northeast) wind that happens immediately after a northeast or a southeast gale, a northwest wind is the only kind that will bring heavy substances ashore, it seems to make the surface current offshore, and this creates an under current on-shore." I have no doubt that *Fulgur perversum* at the locality is raked out of

a fossil bed a short distance offshore, and that this off-shore wind after the on-shore gales favors the tides and currents in doing so.

LEWIS WOOLMAN.

#### THE DEFINITION OF SPECIES.

I HAVE stated in this JOURNAL (N. S., VI, 329) that I believe the quantitative study of variation to be the most pressing problem of biological science. I have consequently read with great interest the papers by Professor Davenport and Mr. Blankinship, on 'A Precise Criterion of Species' (page 685 above). It seems evident that for the definition of species we should not depend on a 'type specimen,' the one first found, in the best state of preservation or the like, but should collate a considerable number of specimens taken at random, and when the traits can be measured give the averages and the mean deviations. Then, as Mr. Davenport explains, we have double-humped curves showing a tendency for the type to split up, and these are of the greatest possible interest to the student of the causes of the evolution of species.

When, however, Mr. Davenport proposes to use a given relation between the height of the smaller hump and the depression between the humps\*—namely 100:50—as a precise criterion

\* This relation depends not only on the distance between the apices, but also on the relative number of specimens of the two types, which, of course, has nothing to do with the difference between the types. There are other cases in Mr. Davenport's paper where the statements seem scarcely to take account of the complexity of the problems. It is meaningless to say that 'in some cases fifty per cent. or even more of the individuals will occur at the mode' and that in this case the curve is steep. The number of individuals at the mode depends on the unit of measurement selected, and the steepness of the curve is arbitrary. The 'half range,' defined as three times the 'standard deviation' (error of mean square), is a theoretically impossible point, and could only be determined approximately from thousands of specimens. Thus in Mr. Davenport's Fig. 9 the 'half range' of the right-hand curve is tripled by a single specimen. In all these cases Mr. Davenport neglects the probable errors which when reckoned show that his distinctions between species and varieties have no validity whatever. The data of Fig. 9 can be expressed by a curve with a single apex.